International Journal of Sciences & Applied Research

www.ijsar.in

METHOD DEVELOPMENT AND VALIDATION BY RP-HPLC - A BRIEF REVIEW

ASHOK GORJA[1], RACHAPUDI GEETHA RANI SAI SUPRAJA*[2], KOYYANA SHIRISHA

[1]Assistant Professor, Department Of Pharmaceutical Analysis, Gokaraju Rangaraju College of Pharmacy, Osmania University Hyderabad-500090, India
[2]research Scholar, Department Of Pharmaceutical Analysis, Gokaraju Rangaraju College of Pharamcy, Hyderabad, India
Corresponding Author: Geetharani.Rachapudi@Gmail.Com

ABSTRACT

Portable stage conditions have a fundamental impact on chromatographic divisions. For instance, before choosing suitable chromatographic conditions, a large number of preparatory trials were conducted using specific mixes of various natural solvents, structures, to obtain acceptable maintenance component, determination, and other chromatographic parameters. The configuration, enhancement, institutionalization, and quality control of pharmaceutical products are made possible by expository procedures. They are equally important for investigations on the medicine digestive system and pharmacokinetics. Accepting systematic systems improvement is necessary to provide administrative entries with reliable data. These techniques are essential for a number of reasons, such as testing for solidity, reference material, quality control discharge, and providing data to support decisions.

INTRODUCTION

One of the most often used methods in pharmaceutical research is UV-visible spectrophotometry. It involves calculating the amount of visible or brilliant radiation that a chemical in an arrangement consumes[1].While there are certain advantages to the HPLC process, most of them involve elevated temperatures, prolonged use of relatively expensive chemicals, extraction, and support structure utilization[2]. In natural chemistry and scientific research, high-performance liquid chromatography (HPLC) is a chromatographic technology that is used to separate mixtures and quantify and refine specific mixture components[3].

In scientific science, high liquid chromatography (HPLC; formerly known as high-weight liquid chromatography) is a technique used to separate the components of a mixture, identify each component, and quantify each component. Pumps are used to move a pressured liquid dissolvable that contains the example combination through a section that has been filled with a potent adsorbent substance[4]. Each component in the example works in slightly different ways with the adsorbent material, resulting in different stream rates for the various segments and causing the components to separate as they flow out of the section[5].

The area of science known as analytical chemistry employs cutting-edge technologies to determine composition through analytical techniques. It is possible to obtain both quantitative and qualitative outcomes[6].It was discovered that the spectrophotometric approach was less

sensitive than the TLC and HPLC techniques. A study indicated that the HPLC approach was more sensitive than the TLC method[7].

Operational pressures in HPLC are substantially greater (50–350 bar), in contrast to regular liquid chromatography, which usually depends on gravity to move the mobile phase through the column. This sets HPLC apart from traditional ("low pressure") liquid chromatography. 2.1–4.6 mm in diameter and 30–250 mm in length are common column dimensions in analytical HPLC because of the minimal sample amount separated. Additionally, the sorbent particles used in HPLC columns are smaller, with an average size of 2 to 50 micrometers. Because of its increased ability to resolve mixtures, HPLC is a widely used chromatographic technique[8].

HPLC procedures for the most part necessitated pricey gear, acquisition for utilization and transfer of solvents, labor focussed specimen arrangement process and person talented in chromatographic strategies. What's more, the great majority of the HPLC routines explored have the prospective application to clinical examination of medication blend, multi-drug pharmacokinetics investigations and collaborative studies[9]

Phase reversal A non-polar stationary phase and a watery, tolerably polar portable phase make up HPLC (RP-HPLC)[10]. Silica that has been surface-adjusted with RMe2SiCl, where R is a straight chain alkyl gathering such as C18H37 or C8H17, is one example of a basic stationary phase. Polar particles elute more quickly (ahead of schedule in the examination) with such stationary phases, whereas less polar atoms require a longer maintenance period.By increasing the amount of water in the portable phase, a specialist can lengthen the maintenance durations. This will strengthen the hydrophobic analyte's natural propensity toward the hydrophobic stationary phase relative to the now-more hydrophilic versatile phase [19]. By increasing the amount of natural dissolvable in the eluent, an examiner can thereby shorten the maintenance period. Because RP-HPLC is used so frequently, it is frequently mistakenly referred to as "HPLC" without additional information. Before releasing sedates, the pharmaceutical industry regularly uses RP-HPLC to evaluate them[11].

Many drugs can be measured with a delicate converse stage ultrafast liquid chromatography (RP-UFLC), which has been developed and approved[8]. The RP-HPLC approach is delicate, repeatable, specific, exact, and particular[12]. The system's main features are its quick investigation time, simple portable stage, easy specimen planning, and improved affectability[13]. Because of this, the method can be used for regular testing in quality-control laboratories[14].

Different types of RP-HPLC segments with unique hydrophobic coatings and pore diameters are available, allowing the expert to regulate the analyte-stationary stage association quality. Generally, longer alkane chains are used for larger analytes, like proteins, and shorter alkane chains are used for peptides and smaller particles to increase the communication quality. Additionally, the example tying qualities and analyte maintenance times can be adjusted by adjusting the steepness of the natural dissolvable angle in the portable stage[15].

Quality control research centers use several investigative systems (HPLC, UV-spectroscopy, HPTLC, Titration, Fluorescence spectroscopy) to ensure the quality, purity, potency, and performance of pharmaceutical goods. Due to a few beneficial circumstances such as speed, specificity, precision, exactness, and ease of mechanization, the HPLC system can analyze the great majority of pharmaceuticals in multicomponent dose structures[16].

The suggested procedure has the advantages of a simple test planning methodology and a brief research duration. The RP-HPLC technology may be used to successfully seek ordinary quantitative estimation and is beneficial in routine lab research with a high degree of exactness and precision. Pharmaceutical companies place a great deal of emphasis on test strategies for all medications, and it is always enticing to select and develop a simple, least time-consuming, accurate, conservative method for determining the medications in API pharmaceutical measurement structures and neurotic specimens like blood and serum [17].

In order to ensure the integrity and quality of the health information supporting investigational new medication applications (INDs), new medication applications (NDAs), abridged new medication applications (ANDAs), and bioavailability (BA), bioequivalence (BE), and bioavailability (BA) studies requiring pharmacokinetic (PK) assessment, agreement with great lab practices (GLPs) for conveying test examination of nonclinical (also known as preclinical) research facility studies and clinical studies is planned. Ensuring that the bioanalytical strategies used are proven effective through advancement, approval (in accordance with a convention, study arrangement, or standard working method [SOP]), and connected as written to effectively evaluate the analyte in the vicinity of a specific network is a definitive goal for any study test examination or system acceptance, regardless of whether GLP agreeability is implemented[18].

A pharmaceutical improvement program's advancement and approval of the expository strategy are essential elements[19]. The HPLC analysis approach is designed to identify, quantify, or sanitize premium mixtures. This technical brief will focus on approval and improvement processes related to pharmaceutical goods. Determining the chromatographic tops for dynamic medicine adjustments was the aim of the system improvement[20].

Recent developments in mass spectrometers and ionization techniques have produced useful new approaches for the study of the medication digestive system and mien. The lack of a sensitive, all-inclusive HPLC finder has led to a surge in interest in high-performance liquid chromatography/mass spectrometry (HPLC/MS)[15]. Although HPLC/MS is not a perfect indication, it has become a viable approach for xenobiotic analysis. The three areas where HPLC/MS is most profitable for investigating the pharmacology and toxicology of atoms with masses less than 1,500 daltons are: enhancing specific routines for follow-up, characterizing and identifying metabolites, and examining interactions between drug particles and peptides/proteins[21].

Strong system development ensures that research center resources are improved, and procedures reach the goals required at each stage of pharmaceutical development[22].Test readiness (percentage natural, pH, shaking/sonication, test size, example age); HPLC investigation conditions (percentage organic, pH, stream rate, temperature, wavelength, and section age); and institutionalization (percentage incorporation, wavelength, standard fixation, and reaction variable adjustment) are the three distinguishing components of an HPLC technique[23].

Prior to the final system improvement, each and every component should be studied during the preliminary strategy advancement plan [46]. Preliminary chromatographic detachment is applied to the debased medicine tests obtained in order to examine the quantity and types of corruption items formed under various circumstances[24].

By altering the pH, which will cause the ionizable analytes to simply separate from the non-ionized structure, maintenance time can be extended[25]. The segment effectiveness can also be increased by adjusting the pH of the flexible stage because this modifies the ionization of the analyte and the residual silanols. It also reduces the possibility of optional communications between the analytes and the silica surface, which could result in an unsatisfactory crest shape[26]. The pH of the portable stage must be changed in order to achieve the optimal determination. The development of the technique can be continued by analyzing the chromatographic partitions' properties at low pH and then at increasing pH until the desired outcomes are obtained[27].

The majority of current HPLC columns favor the use of silica-based packing because of a number of physical properties. For the majority of HPLC separations, fully porous silica particles with a 5 µm diameter offer the required properties[28].

It is possible to improve sample separation by choosing the ideal column temperature. By lowering the viscosity of the mobile phase, a higher column temperature lowers system backpressure and permits the use of longer, more efficient separation columns. The pH at which the quantities of medicines in their ionized and unionized forms are equivalent is known as the acid dissociation constant (pKa). It is a crucial factor in the search for new drugs, especially in physiological systems where the ionization state influences how quickly a substance diffuses across membranes, such as the blood-brain barrier[29].

With the development of HILIC innovation, which has shown to improve repeatability in separating polar and hydrophilic mixes, such as peptides, sugar, vitamins, polar pharmaceuticals, and metabolites, ordinary stage HPLC has recently gained popularity again .In order to effectively expand an HPLC system, the majority of effort should be directed toward method advancement and enhancement, as this will boost the final strategy's execution[30].

VALIDATION

The International Organization for Standardization (ISO) defines confirmation as "procurement of target proof that a given thing satisfies determined necessities", while validation is defined as "check, where the predetermined prerequisites are satisfactory for a proposed utilization". Linearity, accuracy, exactness, toughness, heartiness, LOD, LOQ, and selectivity or specificity are among the various approval parameters [31].

LINEARITY

The ability of a systematic technique to obtain test results that are precisely correlated with the analyte's convergence in the specimen, within a given range, is known as linearity.

A sequence of three to six infusions of five or more norms determines linearity. The adjustment measures' upper ranges, also known as statures, are often plotted in the Y-pivot in relation to the purported standard fixation. The linearity of the shown bend is evaluated by estimating the co-connection coefficient (r2)[32]. The identifier linearity should be estimated using clean standard arrangements, and the strategy linearity should be determined concurrently with the exactness study. The two established criteria for recognizing linearity are: 1) the y-capture should not deviate significantly from zero; and 2) the connection coefficient of the direct relapse line should not be greater than a value that is close to 1[33].

ACCURACY

The degree of agreement between the value that is recognized as a recognized reference quality or as an average real quality and the quality that is found is known as accuracy[34].

PRECISION

It conveys the degree of closeness of understanding (diffuseness) between a series of estimates obtained from different tests conducted on the same homogeneous material in accordance with the suggested parameters. Three criteria can be used to assess accuracy: reproducibility, transitional exactness, and repeatability[35].

Repeatability conveys the accuracy over a brief period of time while using the same working conditions. Another name for repeatability is intra-measure exactness[36].

Within research institutions, intermediate precision communicates in a variety of ways: different days, different investigators, different gear, and so on[37].

Reproducibility is the means by which research centers (whose focus is typically associated with the institutionalization of technology) can communicate accuracy[38].

RUGGEDNESS

An analytical procedure's robustness is a gauge of how well it can withstand slight but intentional changes in method parameters and shows how reliable it is under typical operating conditions[34].

RANGE

The range, for which it has been demonstrated that the systematic approach has an appropriate degree of exactness, precision, and linearity, is the space between the upper and lower convergence of the analyte in the example (counting these foci)[39].

DETECTION LIMIT

The smallest measure of an analyte in an example that can be identified but not precisely quantified as an exact quality is called the detection limit of a particular explanatory approach[40].

QUANTITATION LIMIT

The lowest analyte measure in an example that can be determined quantitatively with appropriate accuracy and exactness is the quantitation limit of a single expository system. A feature of quantitative tests for low mix levels in test lattices, the quantitation limit is particularly useful for identifying contaminating impacts and/or corrupted items[41].

SPECIFICITY

Sometimes, in strategy approval, selectivity and specificity are used in opposition to each other to represent the same idea. The ability to assess the analyte with certainty in the vicinity of components that might be needed as exhibits is known as specificity. Comparing test findings from an analysis of tests with contaminations, debasement items, or placebo fixes with those from an analysis of tests without debasements, corruption items, or placebo fixings regulates the specificity of a test system[42].

REFERENCES

1. Tyagi A, Sharma N et al. HPTLC-Densitometric and RP-HPLC Method Development and Validation for Determination of Salbutamol Sulphate, Bromhexine Hydrochloride and Etofylline in Tablet Dosage Forms. Pharm Anal Acta. 2015; 6: 350.
2. Pavan Kumar C and Gurupadayya BM Analytical Method Development and Validation of Dimethoate Pesticide using HPLC Method. Biochem Anal Biochem. 2013; 2: 127.
3. Atanu Kumar J HPLC: Highly Accessible Instrument in Pharmaceutical Industry for Effective Method Development. Pharm Anal Acta.2012; 3:147.
4. Ankit Tyagi et.al. HPTLC-Densitometric and RP-HPLC Method Development and Validation for Determination of Salbutamol Sulphate, Bromhexine Hydrochloride and Etofylline in Tablet Dosage Forms. Pharmaceut Reg Affairs 2015, 6: 350.
5. Xinxin Zhang et.al. Comparative Studies on Performance of CCC and Preparative RP-HPLC in Separation and Purification of Steroid Saponins from Dioscorea ZingiberensisC.H.Wright. J Steroids Hormon Sci. 2015; 6: 150
6. Chauhan A et.al. Analytical Method Development and Validation: A Concise Review. J Anal Bioanal Tech. 2012; 6: 233
7. Ranjit Singh HPLC method development and validation- an overview. J Pharm Educ Res. 2012 4: 26-33.
8. Yingnian Lu et.al. Development and optimization of a RP-HPLC method to Quantify Midazolam in Rat Plasma after Transdermal Administration: Validation and Application in Pharmacokinetic Study. Pharm Anal Acta 2015, 6: 329
9. Jenkinson C et al. LC-MS/MS-Based Assay for Free and Deconjugated Testosterone and Epitestosterone in Rat Urine and Serum. J Anal Bioanal Tech. 2014; S5: 006.
10. Albert K et.al. Improving the Understanding of the Properties and Retention Behavior of Chemically Bonded Stationary Phases Employing Suspended-state HR/MAS NMR Spectroscopy. J Anal Bioanal Tech.2013; S12: 001.
11. Gengaihi SEl et.al. Antioxidant Activity of Phenolic Compounds from Different Grape Wastes. J Food Process Technol.2014; 5:296
12. Mohammad Faisal Hossain et.al. UV-Metric, pH-Metric and RP-HPLC Methods to Evaluate the Multiple pKa Values of a Polyprotic Basic Novel Antimalarial Drug Lead, CyclenBisquinoline. Mod Chem Appl 2014, 2: 145
13. Najma Sultana et.al. Development and Validation for the Simultaneous Quantification of Prazosin, Amlodipine, Diltiazem and Verapamil in API, Dosage Formulation and Human Serum by RP-HPLC: Application to in vitro Interaction Studies. Med chem 2014, 4: 770
14. Naveed S An Overview of Analytical Determination of Captopril in Active Pharmaceutical Ingredients (API) Formulation and Biological Fluids. J BioequivAvailab. 2013; 5:264-266.
15.Abdallah NA HPLC and Densitometric TLC Methods for Simultaneous Determination of Gemifloxacin with Some Co-administered Drugs in Human Plasma. J Chromatograph SeparatTechniq. 2014; 5:220
16. Vijaya Bhaskar V et.al. Liquid Chromatography/Tandem Mass Spectrometry Method for Quantitative Estimation of Cremophor EL and its Applications. J Anal Bioanal Tech.2013; 4:163.
17. Naveed S Analytical Determination of Lisinopril Using UV Spectrophotometer and HPLC: An Overview. Mod Chem appl.2014; 2: 137.
18. Whitmire M et.al. LC-MS/MS Bioanalysis Method Development, Validation, and Sample Analysis: Points to Consider When Conducting Nonclinical and Clinical Studies in Accordance with Current Regulatory Guidances. J Anal Bioanal Techniques.2011; S4:001.

19. Nouruddin W Ali et.al. Development and Validation of Different Chromatographic Methods for Determination of Two Hypouricemic Drugs in Their Combined Dosage Form. J Anal Bioanal Tech 2014, 5:211
20. Ravi PR et.al. Validation of a Simple, Rapid and Sensitive LC Method for Quantification of Riluzole in Rat Plasma and its Pharmacokinetic Application. J Bioanal Biomed.2012; S6: 007
21. Lories IB et.al. High Performance Liquid Chromatography, TLC Densitometry, First-derivative and First-derivative ratio spectrophotometry for de-termination of Rivaroxaban and its alkaline Degradates in Bulk Powder and its Tablets. J Chromatograph Separat Techniq.2013; 4: 202
22. Gurupadayya BM and Disha NS Stability Indicating HPLC Method for the Simultaneous Determination of Ceftriaxone and Vancomycin in Pharmaceutical Formulation. J Chromat Separation Techniq 2013, 4: 207
23. Apoorva V Rudraraju et.al. In vitro Metabolic Stability Study of New Cyclen Based Antimalarial Drug Leads Using RP-HPLC and LC-MS/MS. Mod Chem Appl 2014, 2: 129
24. Paranthaman R and Kumaravel S A Reversed-Phase High-Performance Liquid Chromatography (RP-HPLC) Determination of Pesticide Residues in Tender Coconut Water (elaneer/nariyal pani). J Chromat Separation Techniq 2013, 4: 208
25. Lories IB et.al. High Performance Liquid Chromatography, TLC Densitometry, Firstderivative and First-derivative ratio spectrophotometry for de-termination of Rivaroxaban and its alkaline Degradates in Bulk Powder and its Tablets. J Chromat Separation Techniq 2013, 4: 202
26. Nada S Abdelwahab et.al. Determination of Thiomersal, Lidocaine and Phenylepherine in their Ternary Mixture. J Chromat Separation Techniq 2013, 4: 199
27.Muhammad Saeed Arayne and Shabana Naz Shah Liquid Chromatographic Analysis of Prazosin in API, Dosage Form and Serum: Application to Drug-Metal Interaction Studies. J Chromat Separation Techniq 2013, 4: 197
28. Anil Chandewar et.al. Method Development and Validation for Desogestrel and Ethinylestradiol in Combined Pharmaceutical Dosage Form by RP-HPLC. Pharmaceut Anal Acta 2013; 4: 262
29. Aparna Gupta et.al. ACE-Inhibitory Activity of Cheddar Cheeses Made with Adjunct Cultures at Different Stages of Ripening. J Adv Dairy Res 2013, 1:102
30. Anandkimar R et.al. Method Development and Validation of Metformine, Pioglitazone and Glibenclamide in Tablet Dosage Form by using RP-HPLC. Biochem Anal Biochem 2013, 2:130
31. Bhatt KK et.al. Simultaneous Estimation of Pregabalin and Methylcobalamine in Pharmaceutical Formulation by RP-HPLC Method. J Anal Bioanal Tech 2012, 4:159
32. Gnana Raja M et.al. Simultaneous, Stability Indicating Method Development and Validation for Related Compounds of Ibuprofen and Paracetamol Tablets by RP-HPLC Method. J Chromat Separation Techniq 2012, 3:155
33. Mehul M. Patel and Disha D Patel Simultaneous Estimation of Metoprolol Succinate and Olmesartan Medoxomil in Pharmaceutical Dosage Form by RP-HPLC. J Chromat Separation Techniq 2012, 3:151
34. Paranthaman R et.al. GC-MS Analysis of Phytochemicals and Simultaneous Determination of Flavonoids in Amaranthus caudatus (Sirukeerai) by RP-HPLC. J Anal Bioanal Techniques 2012, 3:147
35. Jaya Prasanthi K and Syama Sundar B Development and Validation of an HPLC Method for Quantifying Dapiprazole in Bulk Preparations. J Anal Bioanal Techniques 2012, 3:143

36. Kumar P, Ghosh A and Chaudhary M tability Indicating Method Development for Simultaneous Estimation of Ezetimibe and Atorvastatin in Pharmaceutical Formulations by RP-HPLC. Pharm Anal Acta 2012, 3:164
37. Seemi Siddiqui et.al. RP-HPLC Method for Estimation and Stress Degradation Study of Paclitaxel as per ICH Guidelines. J Chromat Separation Techniq 2012, 3:135
38. Jagadeeswaran M et.al. Quantitative Estimation of Lopinavir and Ritonavir in Tablets by RP-HPLC Method. Pharm Anal Acta 2012, 3:160
39. Najma Sultana et.al. RP-HPLC Method for Simultaneous Determination of Captopril and Diuretics: Application in Pharmaceutical Dosage Forms and Human Serum. J Chromatograph SeparatTechniq 2011, 2: 109
40. Subramanian Natesan et.al. Improved Rp- Hplc Method for the Simultaneous Estimation of Tranexamic Acid and Mefenamic Acid in Tablet Dosage Form. Pharm Anal Acta 2011, 2:115
41. Laxman Sawant et.al. Quantitative HPLC Analysis of Ascorbic Acid and Gallic Acid in Phyllanthus Emblica. J Anal Bioanal Techniques 2010, 1:111
42. Nia Y et.al. Determination of Ti from TiO2 Nanoparticles in Biological Materials by Different ICP MS Instruments: Method Validation and Applications. J Nanomed Nanotechnol.2015; 6:269.